



Original Research Article**DOI - 10.26479/2015.0101.02****COMPARATIVE STUDY OF ICHTHYOFAUNA AND FISHERY POTENTIAL OF PERENNIAL TANKS OF KOLHAPUR DISTRICT, (MAHARASHTRA)****S. A. Manjare**

Department of Zoology, Jaysingpur College, Jaysingpur. (M.S), India

ABSTRACT :

Ichthyofauna refers to variety of fish species. The Ichthyofauna community of the perennial tanks was studied by monthly, samples taken from June 2009 to Dec. 2010 in relation to fishery potential. The water bodies such as Tamdalage, Laxmiwadi, Vadgaon tanks were selected. These Tanks are situated at 16⁰ 46' 35' 05" N latitude, 74⁰ 27'48.71 E longitude, 16⁰ 47'23 19" N latitude, 74⁰ 22'56.44" 11 E longitude and 16⁰ 49.45" N latitude, 74⁰ 18'22.50" E longitude respectively, near vicinity of Hatkanangle Tahsil of Kolhapur district, Maharashtra. The water of the tank is used for domestic, Agriculture and fishery activities. Fish collections were done with net of several mesh sizes. 12 fish fauna identified during the study belongs to family Cyprinidae 7 species, Siluridae 1 species, Bagaridae 01 species, and Channidae 2 species of Cichlidae 1 species. Besides identification, relative occurrence and economic importance, fishery potential of fishes of tanks are discussed. All fishes are useful as food fishes expect *Puntius saransa* & *Puntius sophore*. Which are useful as weed and larvicidal fishes . The fish diversity and fishery potential is discussed in detail. To develop a sustainable fishery practices and increase fishery potential of these Tanks.

Keywords – Fish fauna, Abundance, Fishery potential, perennial tanks.

***Corresponding author: S. A. Manjare**

Department of Zoology, Jaysingpur College, Jaysingpur. (M.S), India

Email address: manjaresir@gmail.com

1.INTRODUCTION

Fresh water bodies present a good opportunity for studying the effect of scale on the relative importance of factors that determine diversity on a broad scale. Fresh water bodies are recent and their communities are a combination of species from the former fresh water bodies fish fauna as well as introduced species (Fernando and Holick 1991, Oliveira et. al. 2004) India is one of the mega biodiversity countries in the world and occupies the ninth position in terms of fresh water mega biodiversity as well as mangroves fauna (Mittermeier and Mittermeier,1997 ,Thakar and Sonawane 2013) The Indian fish population represents 11.72 % of species, 23.96% of genera, 57% of families and 80% of the global fishes. Out of the 2200 species so far listed, 73 (3.32%) belong to the cold freshwater regime 544 (24.73%) to the warm fresh waters domain, 143(6.50%) to the brackish waters and 1440 (65.45%) to the marine ecosystem) (I CBD1994).The Indian fish fauna is divided into two classes, viz Condrichthyes and Osteichthyes. The Condrichthyes are represented by 131 species under 67 genera, 28 families and 10 orders in the Indian region. The Indian Osteichthyes are represented by 2,415 species belonging to 902 genera, 226 families and 30 orders of which five families, notably the family Parapsilornychidae are endemic to India. The country is endowed with vast and varied resources possessing river ecological heritage and rich biodiversity. Freshwater fishery sites are varied like 45,000 km of rivers, 1,26,334 km of canals, ponds and tanks 2.36 million ha, 2.05 million ha of reservoirs (Ayyapan, 2004, Thakar & Sonawane 2013) India's inland water resources are diversified, as they are plentiful. Reservoirs contribute the single largest inland fishery resources both in terms of size & production potential. Fish fauna of a reservoir basically represents the fish diversity and their abundance. Indian reservoir preserve a rich variety of fish species, which supports to the commercial fisheries (Kantaraj et. al 2011).Fresh water bodies in Kolhapur district of Maharashtra state have not been studied in detail except the sporadic work of Sawant (1978), Angadi (1985) Gaikwad (1996), Pailwan (2005) etc. Their Contribution in the field of limnology and hydrobiology of this area serve as a guideline to the new researchers of this region. Therefore present investigation is formulated to assemble the information regarding ichthyofauna and fisheries potential of these tanks namely Tamdalage, Laxmiwadi and Vadgaon of Kolhapur district of Maharashtra state.

2.MATERIALS AND METHODS

Study Area

The Tamdalage, Laxmiwadi and Vadgaon tanks are perennial freshwater tanks located $16^{\circ} 46' 35''$ N latitude, $74^{\circ} 27'48.71$ E longitude, $16^{\circ} 47'23 19''$ N latitude, $74^{\circ} 22'56.44''$ 11 E longitude and $16^{\circ} 49.45''$ N latitude, $74^{\circ} 18'22.50''$ E longitude respectively, in the vicinity of Hatkanangle Tashil. These tanks are basically used for drinking water, for domestic purpose and fishery activities by local fisherman communities. These tanks are auctioned on lease for the period of 3-5 years by Zilla Parishad Kolhapur. These tanks are situated in hilly region. They have only rain water source. These tanks exhibit more fluctuations in its water level and characterized by more anthropogenic activities & silty bottom and huge algal bottom throughout year. One the most characteristic feature of these tanks is absence of molluscan fauna. The length of the wall is 574.12 & 875.86 meters its height is about 23, 20 & 31 feet. They are located at an elevation of 2066, 2090 & 2089 feet above mean sea level. The fishes were collected from these tanks with help of local fisherman monthly basis during the year Jan 2009 to Dec. 2010. The fishes were preserved in 10% formaldehyde Solution for taxonomic analysis. Identification and Economic importance of fishes were carried out with the help of standard literature (Day 1958, Datta & Shrivastara 1988, Jhingran 1991 & Jayaram1981) for fishery potential frequent visits were made to attend the commercial catches of these tanks were analyzed in order to obtain information regarding fish species, minimum and maximum length & weight species wise, total weight & dominant species in the catches. The approximate annual productivity were calculated considering 200 operations per year excluding off season & closed period. The fishery potential status of these tanks were estimated based on information regarding the capital and operational investments, approximate annual fish production wholesale rates /kg of fish. The fish production per acre was estimated considering the average water spread area in hectares, multiplied by 2.5 provides an area gives fish production per acre. This is turn multiplied by wholesale rate of Rs. 60 per Kg of fish resulted in to per acre income from fish production of these tanks.

3. RESULTS AND DISCUSSION

The Ichthyofauna is mainly depends upon the biotic and biotic factors and types of the ecosystem, age of water bodies, depth, water level fluctuations, morphometric features and bottom have great implications. The physico-chemical and biological features of the collection centers also play an effective role in fisheries out put to a greater extent. Among 11 species of fishes the family Cyprinidae was most dominant in the assemblage composition with 68.25% followed by Cichlidae 21.45% with Channidae 6.12 and Bagaridae and Silurian 4.18% respectively. The present-results get support from other workers like Kantaraj et al (2011) Venkatswarlu et. al (2007) Wakid and Biswas (2005). The monthly commercial catches observed at Tamdalage tank the fish species such as *Oriochromis Mossambicus*, *Channa punctatus*, *Puntius sarana*, *Cirrhinus mrigala*, *Labeo rohita*, *Catla catla* and *Cyprinus carpio* caught through out the study period However, Species like *Wallago attu*, *Mystus Seenghala*, *Channa striatus* are moderately observed from the catches. In Laxmiwadi tank predominant of carps species such as *Cirrhinus mrigala*, *Labeo rohita*, *Catla catla* and *Cyprinus carpio*, with infrequent occurrence of predatory fishes such as *Wallago attu*, *Channa striatus* was observed. From commercial catches of Vadgaon tank represented mostly the predominance of introduced species such as *Cirrhinus mrigala*, *Catla catla* and *Cyprinus carpio* with minor contribution of fishes like *Wallago attu*, *Channa striatus*, *Mystus seenghala* and *Puntius sarana*, *Puntius sophore* *Channa punctatus* are less dominant species. Similar pattern of fish diversity was observed by Lagler (1953) classified the fish species on the basis of their economic importance in Bhadra reservoir of Karnataka. Ahirrao and Mane (2000) recorded 52 fish species belongs to 8 families from fresh waters of Parbhani district of Marathwada region of Maharashtra state. Sakhare (2001) identified 23 fish species belonging to 7 order in Jawalgaon reservoir in Solapur district. Pailwan (2005) also reported that 23 fish species belonging to 11 orders from freshwater tanks in Kolhapur district of Maharashtra. The present study of fish fauna in these tanks showed that most of the fish species recorded were widely distributed in various region of Kolhapur district. The fish species like carp groups were more dominant. Therefore the present investigation reveals that Cyprinide fishes are found to be more dominant group than others which is supported by other studies also (Singh et al 2006). From commercial fish catches revealed that, the minimum average length and weight for naturally occurring fish species was (21.09 cm and 196.72 gm), (45.5 cm and 587.5 gm) and (30.48 cm and 220.67 gm) during 2009-2010 at Tamdalage, Laxmiwadi and

Vadgaon tanks respectively. Similarly the average maximum length and weight for naturally occurring was recorded as (31.37 and 421.86 gm), (46.75 cm and 587.5 gm) and (44.82 cm and 722.57 gm) during 2009-2010 at Tamdalge, Laxmiwadi and Vadgaon tanks. From commercial fish catches revealed that, the minimum average length and weight for carp fish species was (52.10 cm and 240.18 gm), (32.12 cm and 831.87 gm) and (31.88 cm and 269.15 gm) during 2009-2010 at Tamdalge, Laxmiwadi and Vadgaon tanks respectively. Similarly the average maximum length and weight for carp fish species was recorded as (67.41cm and 602.03 gm), (47.34 cm and 2074.24 gm) and (46.11 cm and 841.84 gm) during 2009-2010 at Tamdalge, Laxmiwadi and Vadgaon tank. (Table -1). Based on the commercial catches the average mean catch was estimated as 34.88 kg at Tamdalge tank. It was recorded 30.33 kg and 26.50 kg at Laxmiwadi and Vadgaon tank during the year 2009-2010 respectively. However, based on available information, the economic status of these tanks was estimated considering the capital and operational cost invested for fish culture as mentioned in economic status. In Tamdalage tank the capital and operational cost was estimated up to Rs 1, 00, 000, the seasonal commercial catches revealed the approximate fish production up to 6.97 tons and annual income from the sale of fish was estimated up to Rs 4, 18,200 Excluding the cost of production the net profit was estimated to about Rs 3, 18,200 during year 2009 - 2010 respectively (Table- 1).In Laxmiwadi tank the capital and operational cost was estimated up to Rs 70,500. The seasonal commercial catches revealed the approximate fish production up to 6.06 tons and annual income from the sale of fish was estimated up to Rs 3, 63, 600 excluding the cost of production the net profit was estimated to about Rs 2, 93, 400 during year 2009 - 2010 respectively (Table- 1).In Vadgaon tank the capital and operation cost was estimated up to Rs 82, 000 the seasonal commercial catches revealed the approximate fish production up to 5.30 tons and annual income from the sale of fish was estimated up to Rs 3, 18, 000 Excluding the cost of production the net profit was estimated to about Rs 2, 36, 000 during year 2009 - 2010 respectively (Table- 1).Considering the average water spread area, approximate fish production, no. of catches and net income/ acre Tamdalge tank represented maximum fish production, followed by Laxmiwadi & Vadgaon tanks. However, scientific management including proper stocking density, supplementary feeding, manuring and preservation were observed as ill-considered in these tanks.

Sr. No.	Expenditure	Tamdalge	Laxmiwadi	Vadgaon
A	Capital cost	Amount in Rs.	Amount in Rs.	Amount in Rs.
1	Lease amount	40,000	25000	35000
2	Expenditure on fish seed including transportation	20,000	15000	20000
B	Operational cost			
3	Expenditure on nets & gears	5,000	10,000	10,000
4	Cost of harvesting	30,000	15,000	15,000
5	Expenditure on harvesting	3,000	3,000	3,000
6	Miscellaneous	2,000	2500	2000
	Total (A+B)	1,00,000	70,500	82,000
C	Production			
7	Fish production with an average catch 30.87 & 34.88 kg/ operation	6.976 tones	6.066 tones	5.300 tones
8	Total income from the sale of fish with the wholesale rate (Rs.60/kg)	4,18,560	3,63,960	3,18,000
9	Net profit from fish production (A+B-C)	3,18,560	3,41,040	2,36,000
10	Net income per acre	25,625	15,238	10,124

Table 1. Economic status in terms of expenditure, fish production and net profit per acre at Tamdalage, Laxmiwadi & Vadgaon Tanks (Jan2009-Dec2010)

REFERENCES

1. Ayyapan S, Birdar S. R (2004): Enhancing Global Competition, Survey Of Indian Agriculture (The Hindu).
2. Thirumala. S, Kiran B.R, and Kantaraj. G. S (2011): Fish diversity in relation to physico-chemical characteristics of Bhadra reservoir of Karnataka, India. *Advances in Applied Science Research*. 2 (5) : 3447
3. Samant, J. S. (1978): Studies on hydrobiology of Kalamba reservoir, Kolhapur. Ph. D. Thesis University of Bombay, Bombay.
4. Angadi, S. M. (1985): Hydrobiology of Rajaram tank. M. Phil. Dissertation, Shivaji University, Kolhapur.
5. Gaikwad, P. T. (1996): Hydrobiology of Shirolu reservoir. M. Phil. Dissertation, Shivaji University, Kolhapur.
6. Pailwan, I. F. (2005): Limnology and fisheries potential of perennial tanks of Kolhapur District. Ph. D. Thesis Shivaji University, Kolhapur.
7. Day, Francis (1958): The fishes of India Vol. I & II. William Dawson and sons Ltd. London.
8. Datta Munshi, Srivastava M. P (1988): Natural history of fishes and systematics of fresh water fishes of India. Narendra Publishing House, Delhi.
9. Jhingran, A. G. (1982): Fish and fisheries of India 2nd edition. Hindustan Publishing Corporation(India)Delhi.
10. Jayaram K. C (1981): The fresh water fishes of India, Pakistan, Burma and Sri-Lanka. Handbook Zoological Survey of India, No. 2. 475.
11. Venkatshwarlu M, Jothi Srigowri, Asha Shree H.M (2007): Proceeding on Diversity and Life Processes from Ocean and Land". Goa University, Goa. 156-15
12. Sambhaji B. Thakar, Kailas D. Sonawane (2013) Mangrove Infoline Database: A Database of Mangrove Plants with Protein Sequence Information *Current Bioinformatics*. 8.524- 529.
13. Waked A, Biswas S. P (2005): *Journal of Bombay Natural History Society*. 102 (1), 50-55.
14. Lagler K. F (1953): *Freshwater fishery biology* W.M.C. Brown and Co. Iowa.
15. Ahirrao S.D, Mane A.S (2000): *Journal of Aquatic Biology* 15 (1&2), 40-43
16. Sakhare V.B.(2001): *Journal of Aquatic Biology* 16 (1&2), 31-33 Singh et al 2006)
17. APHA (1998): *Standard Methods for the Examination of Water and Wastewater*. 20th edition. Amer. Publ. Health. Assoc., Amer. Water Works Assoc. and Water Poll. Contr. Fed., Washington, D.C.