**Original Research Article**

DOI: 10.26479/2018.0404.31

**BIOREMEDIATION OF LEAD CONTAMINATED SOIL USING BACTERIA**Abhishek Kumar Pandey<sup>1\*</sup>, Abhilasha Shrivastava<sup>2</sup>

1.Center for Biotechnology Studies, Awadhesh Pratap Singh University, Rewa, MP, India.

2. Department Biotechnology, Government Model Science College, Rewa, MP, India.

**ABSTRACT:** Bioremediation is the use of living organisms (primarily microorganisms) for removal of a pollutant from the biosphere. It relies on biological processes to minimize an unwanted environment impact of the pollutants. The microorganisms in particular have the abilities to degrade, detoxify and even accumulate the harmful organic as well as inorganic compounds. Soil samples were collected from J P Cement Plant, Rewa from different places at a depth of 6 -10 Inches from 4 - 6 different spots. These soil samples were mixed properly and enriched for Lead resistant clones by incubating 10g of slag in 90 ml of sterile water amended with 10 ml Luria Bertani (LB) medium and 20 µg/ml each of lead-acetate at 37°C for 2h. Supernatants were plated at 10<sup>-2</sup> dilution by spread-plate method on LB agar medium. The plates were then incubated at 37°C. The colonies appeared after 3 days. Strains were preserved and phenotypic studies were carried out. The metal accumulation efficiency was measured by Atomic Absorption Spectroscopy (AAS). The soil samples collected are all alkaline in nature, the isolated bacteria are gram positive, rod shaped, aerobic, salt tolerant, endospore forming bacteria and according to this research sample with bacterial inoculation in them shown reduction in the Lead levels as compared to the raw soil samples.

**KEYWORDS:** Bioremediation, Heavy metals, Lead, Soil, Bacteria.

**Corresponding Author: Abhishek Kumar Pandey**

Center for Biotechnology Studies, Awadhesh Pratap Singh University, Rewa, MP, India.

Email Address: abhi73025@gmail.com

**1. INTRODUCTION**

Heavy metals have a major problem to human health and environmental issues due to the high incidence as a contaminant, low solubility in biota and classification of various heavy metals as carcinogens and mutagens [1]. Heavy metals can produce harmful effects on human health when

they are taken up in amounts that cannot be processed by the organism. In addition, these metals cannot be degraded to harmless products and hence persist in the environment indefinitely [2]. Contamination of heavy metals in the environment is a major global concern, because of toxicity and threat to the human life and ecosystem [3]. The levels of metals in all environments, including air, water and soil are increasing in some cases to toxic levels, with contributions from wide variety of industrial and domestic sources [4]. Metal contaminated environments pose serious threat to health and ecosystems. Metals like arsenic, cadmium, lead, mercury etc cause conditions including hypophosphatemia, heart disease and liver damage, cancer and neurological and cardiovascular diseases, central nervous system damage and sensory disturbances [5]. Bay incident of Japan is an example of heavy metal poisoning which occurred due to consumption of fishes and shellfishes contaminated with methylmercury in their body [6]. The ecological effects of toxic metals and their biological magnification through the food chain have prompted a demand for decontamination of heavy metals [7]. Lead has been introduced in the environment from a variety of sources such as, storage battery, lead smelting, tetraethyl-lead manufacturing and mining, plating, ammunition, ceramic and glass industries [8]. It has been used in piping, building materials, solders, paint, type metal, ammunition and castings since the medieval times [9]. Lead can cause anemia, kidney disease, harm to nervous system and can also cause wildlife destruction [10]. Several methods have been designed for the treatment and removal of heavy metals in contaminated site. Physico-chemical methods have been used, such as electrochemical treatment, ion exchange, precipitation, reverse osmosis, evaporation, and sorption [11, 12]. Ex situ remediation techniques involve removing the soil from the subsurface to treat it. Insitu remediation techniques involve leaving the soil in its original place and bringing the biological mechanisms to the soil. But these methods have disadvantages, including economically expensive, incomplete metal removal, requirements higher reagent energy, and generation of toxic sludge [13, 14]. Because of toxicity and the ubiquity of the metals in environment, microbes have developed unique and sometimes bizarre ways of dealing with unwanted metals. Some microorganisms have mechanisms to sequester and immobilize metals, whereas others actually enhance metal solubility in the environment. Sometimes they oxidize or reduce them to a non-toxic or relatively less toxic forms [15]. Bioremediation is a sustainable strategy that utilizes the metabolic potential of microorganisms and plants to clean-up contaminated environments [16]. It achieves contaminant decomposition or immobilization by exploiting the existing metabolic potential of microorganisms with novel catabolic functions derived from selection or by introduction of genes encoding such functions [17]. The most essential parameters required for bioremediation are the nature of pollutants, soil structure, temperature, pH, moisture content, hydrogeology, the nutritional state, redox-potential, and microbial diversity of the site. In bioremediation processes, microorganisms use the contaminants as nutrient or energy sources [18]. Bioremediation is a cost effective eco-friendly means of healing nature with nature. This technology

Pandey & Shrivastava RJLBPCS 2018 www.rjlbpcs.com Life Science Informatics Publications  
may be applied in the removal of xenobiotic compounds from agrochemical and petrochemical industries, oil spills, heavy metals in sewage, sludge and marine sediments etc [19].

## **2. MATERIALS AND METHODS**

### **2.1 Collection of soil samples**

Soil samples were collected from J P Cement Plant, Rewa from different places at a depth of 6 -10 Inches from 4-6 different spots.

### **2.2 Physicochemical characteristics**

Different physico-chemical characteristics of the leachate such as pH, Dissolved Oxygen (DO), alkalinity and total water content were determined for both freshly produced slag and old slag deposited considerably since long period of time. Methodologies followed were according to American Public Health Association (APHA) [20].

### **2.3 Strains isolation**

The samples were mixed properly and enriched for lead resistant clones by incubating 10g of slag in 90 ml of sterile water amended with 10ml Luria Bertani (LB) medium and 20 µg/ml of lead-acetate at 37 °C for 2h. Supernatants were plated at 10<sup>-2</sup> dilution by spread-plate method on LB agar medium containing 20 µg/ml of lead-acetate [Pb (CH<sub>3</sub>COO)<sub>2</sub>]. The plates were then incubated at 37 °C. The colonies which appeared after 3 days of incubation were further screened at higher concentrations (20-800 µg/ml) of heavy metal. Finally, the strains were selected as lead resistant isolates for further studies.

### **2.4 Maintenance and preservation of cultures**

Strains were preserved in the refrigerator in stabcultures made of LB medium and NA medium both for short time preservation. The media were all the time supplemented with 20 µg/ml of heavy metal and for long time preservation glycerol stocks were made for storage in – 70 °C for longer period of time. For this overnight grown liquid culture were taken in cryo-vials and added with 15% glycerol after this the vials were transferred to – 70 °C deep freezer.

### **2.5 Phenotypic studies and Determination of Maximum Tolerance Limit (MTL)**


Phenotypic studies such as colony morphology, Gram staining, endospore staining, motility, biochemical tests, acid production tests etc. were carried out as per standard methods of Benson, 1990 in our laboratory [21]. MTL was determined by growing cells both in LB broth and LB agar media which had been amended with increasing concentrations of respective heavy metal. Heavy metal (Pb) stock were prepared in sterile distilled water and was added at the time of inoculation of specific bacterial isolates in inoculating chamber.

### **2.6 Heavy metal accumulation efficiency measurement**

The heavy metal accumulation efficiency was measured by Atomic Absorption Spectroscopy (AAS). For this 500 mg cell pellets obtained at different time intervals were suspended in 20 ml water added with 5% conc. HNO<sub>3</sub> and 0.5% conc. HCl. The cell suspensions were then digested in Anton Paar

### 3. RESULTS AND DISCUSSION

The physico-chemical characteristics indicate that the place is unsuitable for the growth of plant mainly because of high pH. The older slag is more alkaline than the slag deposited recently. The dissolved oxygen of the older slag is also considerably higher than the fresh slag. The lead resistant strain is a gram positive, rod shaped, aerobic, salt tolerant, endospore forming bacteria. To study the maximum tolerance limit (MTL), cells were grown in LB broth with increasing concentrations of heavy metals and growth was observed, the maximum tolerance limit of lead resistant bacteria was 900 µg/ml. The concentrations of heavy metal were determined using Atomic Absorption Spectroscopy (AAS) as described in materials and methods. The concentrations were determined at 12 hrs, 18 hrs, 24 hrs and 48 hrs of growth. it can be seen that the accumulation of heavy metal (lead) increases gradually in log phase and they show maximum level of accumulation at late log phase to stationary phase (i.e. at 18-20 hrs), however, the accumulation takes place even after the death of the cells in lead resistant strains, this indicates involvement of a passive mode of heavy metal accumulation. The maximum accumulation efficiencies of lead resistant strain as determined by atomic absorption spectroscopy were 0.723 mg/g (Figure 1).



**Figure 1: Heavy metal accumulation efficiencies at different time intervals**

The principal purpose of this study was to isolate and characterize some heavy metal resistant bacteria from slag with respect to their heavy metal accumulation efficiencies. As has been discussed earlier that after a few rounds of screening at consecutively increasing concentrations of lead, bacterial isolates were screened and selected for further studies. The alkaliphilic nature and salt tolerant abilities of the strains are significant with respect to their application in metal contaminated, salinated, alkaline soil for bioremediation. Moreover, the slag, the natural habitat from which these bacteria were isolated also had a pH of 9.8. The increased metal tolerance in relation to pH has been

Pandey & Shrivastava RJLBPCS 2018 www.rjlbpccs.com Life Science Informatics Publications reported in *Aspergillus* sp [22]. The lead accumulation efficiency of the strain as determined by atomic absorption spectroscopy was found to be optimum at 18 hours after inoculation when the population reached almost the maximum density. The accumulation efficiency then decreased considerably indicating it to be an active uptake system requiring energy for transport across the membrane [23]. It has been shown that microbes have the capability to reduce heavy metals, but whether they reduce it for detoxification or for growth is a matter of concern. In a sample of agricultural soil, arsenate reduction was not coupled to growth [24], rather it could be linked to detoxification. However, in the JMM-4 strain of *Bacillus* sp., a close relative of *Bacillus arsenicoselenatis* isolated from arsenic contaminated mud in Australia had been capable of oxidizing lactate to acetate while reducing arsenate to arsenite [25]. Regarding lead resistance there are reports that level of resistance for lead is also directly related with the biomass which is in accordance with our study [26].

#### **4. CONCLUSION**

Conclusively, the bacterial isolates isolated from the slag disposal site of JP Cement factory Rewa, MP, India have been characterized in the present study with respect to their heavy metal acquisition and resistance and according to this research sample with bacterial inoculation in them shown reduction in the lead levels as compared to the raw soil samples. Thus it has been concluded that bioremediation is a potential method for solving the problem of heavy metal pollution

#### **ACKNOWLEDGEMENT**

The author thankfully acknowledges Dr. Sher Singh Parmar and Dr. Arvind Shukla, faculty members of Center of Biotechnology studies, APS University, Rewa for their usefull and important guidance.

#### **CONFLICT OF INTEREST**

There is no conflict of interest

#### **REFERENCES**

1. Brooks RR. Plants that hyperaccumulate heavy metals. Cambridge University Press, London, UK. 1998; 92-96.
2. Jones CA, Langer HW, Anderson K, McDermott TR, Inskeep WP. Rates of microbially mediated arsenate reduction and solubilization. Soil Science Soc. Am. J. 2000; 64: 600-608.
3. Van Deuren J, Lloyd T, Chhetry S, Raycham L, Peck J. Remediation Technologies Screening Matrix and Reference Guide. Federal Remediation Technologies Roundtable. 2002; 5: 1-4.
4. Koning M, Hupe K, Stegann R. Thermal Processes, Scrubbing/Extraction, Bioremediation and Disposal. Biotechnology. 2000; 11b: 306-317.
5. Xu S, Sheng G, Boyd SA. Use of organoclays in pollutant abatement. Adv. Agron. 1997; 59: 25-62.
6. Schulz-Berendt V. Bioreediation with Heap Technique. Biotechnology. 2000; 11b: 320-328.
7. Kadirvelu K, Thamaraiselvi K, Namasivayam C. "Removal of heavy metals from industrial

- Pandey & Shrivastava RJLBPCS 2018 www.rjlbpcs.com Life Science Informatics Publications  
waste water by adsorption onto activated carbon prepared from agricultural solid waste.”  
Bioresource Technology. 2001; 6: 63 – 65.
8. Held T, Don H. In Situ Remediation. *Biotechnology*. 2000; 11b: 350-370.
  9. Congeevaram S, Dhanarani S, Park J, Dexilin M, Thamaraiselvi K. Biosorption of chromium and nickel by heavy metal resistant fungal and bacterial isolates. *J. Hazard Mater*. 2007; 146: 270-277.
  10. Iranzo M, Sainz-Padro I, Boluda R, Sanchez J, Mormeneo S. The Use of Microorganisms in Environmental Engineering. *Annals of Microbiology*. 2001; 51: 135-143.
  11. Guidi W, Palm E, Mancuso S, Azzarello E. “Trace element phytoextraction from contaminated soil. a case study under Mediterranean climate”. *Environmental Science and Pollution Research*. 2018; 18: 1197-1205.
  12. Xiong J, Wu L, Tu S, Van Nostrand, JD, He Z, Zhou J. Microbial communities and functional genes associated with soil arsenic contamination and the rhizosphere of the arsenic-hyperaccumulating plant. *Environ. Microbiol*. 2010; 76: 7277–7284.
  13. Naz T, Khan MD, Ahmed I, Rehman SU, Rha ES, Malook I. Biosorption of heavy metals by *Pseudomonas* species isolated from sugar industry. *Toxicol. Ind. Health*. 2015; 10: 525-534.
  14. Shannon M.JR, Unterman R. Evaluation Bioremediation: Distinguishing Fact from Fiction. *Annual Review of Microbiology*. 2000; 47: 715-738.
  15. Donald LS. *Environmental Soil Chemistry*. Academic Press. San Diego, California, USA. 2003; 87-92.
  16. Kleijntjens RH, Luyben KC. Bioremediation. *Biotechnology*. 2000; 11b: 329-347.
  17. Pandey S, Saha P, Biswas S, Maiti TK. Characterization of two heavy metal resistant strains isolated from slag disposal site at Burnpur. *Indian J. Environ Biol*. 2011; 32: 773-779.
  18. Faryal R, Sultan A, Tahir F, Ahmed S, Hameed A. Biosorption of lead by indigenous fungal strains. *Pak. J. of Bot*. 2007; 39: 615 – 622.
  19. Zhiqi H, Reiske HR, Wilson DB. Characterization of cadmium uptake in *Lacto plantarum* and isolation of cadmium and manganese uptake system. *Applied and Environmental Microbiology*. 1999; 65: 4741-4745.
  20. American Public Health Association (APHA). *Standard methods for the examination of water and wastewater*. 19<sup>th</sup> American Public Health Association, Washington, DC. 1995; 18: 65 – 73.
  21. Benson HJ. *Microbiological Application – A Laboratory Manual in General Microbiology*. C. Brown Publishers: USA. 1990; 5: 125 - 146.
  22. Konopka A, Zakharova T, Bischoff M, Oliver L, Nakatsu C, Turco RF. Microbial biomass and activity in lead-contaminated soil. *Appl Environ Microbiol*. 1999; 65: 2256-2259.
  23. Rani A, Ray K, Goel R. Strategies for crop improvement in contaminated soils using metal-tolerant bioinoculants. *Springer*. 2010; 11: 105-132.

24. Akhtar MS, Chali B, Azam T. Bioremediation of Arsenic and Lead by Plants and Microbes from Contaminated Soil. *Res. Plant Sciences*. 2013; 1 (3): 68-73.
25. Jain R, Ahmad P, Gadgil K, Sarma S. Heavy metal toxicity: Effect of plant growth, Biochemical parameters and metal accumulation by *Brasica juncea*, L. *Int.J. Plant Production*. 2012; 3 (3): 65-75.
26. Santini JM, Stolz JF, Macy JM. "Isolation of a new arsenate respiring bacterium –physiological and Phylogenetic studies," *Geomicrobiol. J*. 2002; 19: 41-52.