



Original Research Article

DOI:10.26479/2019.0501.45

BIOCHEMICAL CHANGES DUE TO INTEGRATED DISEASE MANAGEMENT APPROACHES FOR BROWN LEAF SPOT DISEASE IN RICE VAR. ADT 36

Jaiganesh V*, Kannan C.

Department of Plant Pathology, Faculty of Agriculture, Annamalai University,
Annamalai nagar, Cuddalore DT, Tamil Nadu, India.

ABSTRACT: The Pot culture studies were undertaken to investigate the changes of amino acid, Protein and starch compound in rice as influenced by application of Zinc sulphate and foliar application of salicylic acid and Potassium silicate and Brown spot pathogen *Bipolaris oryzae* inoculation. The results revealed that Soil application of Zinc sulphate @ 25 kg/ ha along with foliar application of plant activator Salicylic acid @ 50 ppm on 15 days after transplanting and Foliar spray of silicon based nutrient potassium silicate @ 3 % recorded the minimum disease incidence and maximum biometrics of rice. The same treatment recorded the minimum amount of amino nitrogen content whereas protein and starch content increased when compared to comparison fungicide and control treatments. Protein content increased up to the 14th day of sampling and then slightly reduced in all the treatments. The minimum protein content was recorded in control at final sampling.

KEYWORDS: Rice brown spot, Amino nitrogen, Protein, Integrated Disease Management.

Corresponding Author: Dr. Jaiganesh V* Ph.D.

Department of Plant Pathology, Faculty of Agriculture, Annamalai University, Annamalai nagar
Cuddalore DT, Tamil Nadu, India. Email Address: potatojaiganesh@gmail.com

1.INTRODUCTION

Rice (*Oryza sativa* L.) is the second most cultivated crop worldwide and it has been estimated that half the world's population survives wholly or partially on this crop [1] and rice provides more calories per ha than any other cereal food grains. In India 136.5 million tonnes of rice was produced from an area of 44.0 million ha with the productivity of 2915 kg per ha [2]. In Tamil Nadu, rice is cultivated in an area of 2.05 million ha with a production of 7.2 million tonnes [3].

© 2019 Life Science Informatics Publication All rights reserved

Peer review under responsibility of Life Science Informatics Publications

2019 Jan – Feb RJLBPCS 5(1) Page No.540

Rice crop is widely affected by a number of diseases caused by fungi, bacteria, viruses and mycoplasma which results in considerable yield losses [4]. Among the various fungal diseases of rice, brown spot or sesame leaf spot incited by *Helminthosporium oryzae* (Breda de Haan) Subram. and Jain (Current name: *Bipolaris oryzae* (Breda de Haan) Shoemaker) is found to occur in most rice growing areas. Currently the disease is being managed by application of fungicides. Due to pesticides hazards, pollution effect, fungicide resistant, bio control agent resistant strains, lack of bioprotectant knowledge which required the integrated component approach in Indian farmer's level which will be improve growth and disease suppression. The development of disease and pathogen establishment involves chemical interactions between host and pathogen. The host plant reacts to the stimulus of the intruding pathogen which is manifested by the altered metabolism of the plant. Alterations in the host physiology as a result of pathogenic invasion have been well recognized in a number of host parasite interactions. Biochemical reactions will determine the nature and some extent of resistance or susceptibility. The initial response of a plant to infection was presumably induced by a parasite and subsequent alterations were the results of host's re-adjustments to the pathogens [5]. Therefore, with an aim to develop an integrated strategy involving the use of certain macro-micro nutrients, silicon based nutrients and resistance inducing chemicals for the successful sustainable management of rice brown spot. Hence, the present studies were undertaken to investigate the changes of Phenol content by application of Macro-micro nutrient, Salicylic acid, potassium silicate along with pathogen inoculation.

2. MATERIALS AND METHODS

Crop, Variety and Source

Crop : Rice (*Oryza sativa* L.)
Variety : ADT 36
Source : Tamil Nadu Rice Research Institute (TRRI),
Aduthurai, Tamil Nadu.

Pot culture studies

The pot culture studies was conducted to test the efficacy of certain macro-micro nutrients, silicon based nutrients and certain resistance inducing chemicals for assessing their influence on the incidence of brown spot of rice with various treatment and combinations. The brown spot susceptible variety ADT 36 grown in rectangular pots of size, 30x45 cm was used for the study. The plants were given artificial inoculation by spraying the spore suspensions with adequate spore load (50,000 spores/ml) at 15 DAT in the evening hours. The crop was maintained in a poly house with frequent spraying of water to provide adequate moisture and relative humidity to enable successful infection by the pathogen. The experiments were conducted in a randomized block design with three replications for each treatment and a suitable control. The fungicide carbendazim 50 WP @ 0.1 per cent was used for comparison and the standard agronomic practices as

recommended by the State Agricultural Department were followed. The effective treatments observed in different experiments conducted under pot and field conditions were pooled together and a new schedule of treatments in combination was evolved for the effective management of brown spot disease of rice. Also, zinc sulphate @ 25 Kg/ha was applied as basal application to the entire treatments (ZSS) except control and comparison. The treatment details are given below;

Treatment schedule

T₁ – ZSS + ZSF₁ + ZSF₂

T₂ – ZSS + SA₁ + SA₂

T₃ – ZSS + PS₁ + PS₂

T₄ – ZSS + ZSF₁ + SA₂

T₅ – ZSS + SA₁ + ZSF₂

T₆ – ZSS + SA₁ + PS₂

T₇ – ZSS + PS₁ + SA₂

T₈ – ZSS + PS₁ + ZSF₂

T₉ – ZSS + ZSF₁ + PS₂

T₁₀ – Carbendazim 50 WP @ 0.1 per cent as foliar spray (comparison fungicide)

T₁₁ – Control

ZnSO₄ @ 25 Kg/ha was applied as basal application to the entire treatments (ZSS) except control and comparison. The treatment details are given below;

T₁ – ZSS + Two sprays of zinc sulphate @ 3 % on 15 and 30 DAT

T₂ - ZSS + Two sprays with salicylic acid @ 50 ppm on 15 and 30 DAT.

T₃ - ZSS + Two sprays with potassium silicate @ 3 % on 15 and 30 DAT.

T₄ - ZSS + First spray with zinc sulphate @ 3 % on 15 DAT + second spray with salicylic acid @ 50 ppm on 30 DAT.

T₅ - ZSS + Second spray with zinc sulphate @ 3 % on 30 DAT

T₆ - ZSS + First spray with salicylic acid @ 50 ppm on 15 DAT + second spray with potassium silicate @ 3 % on 30 DAT

T₇ - ZSS + First spray with potassium silicate @ 3 % on 15 DAT + second spray with salicylic acid @ 50 ppm on 30 DAT

T₈ - ZSS + First spray with potassium silicate @ 3 % on 15 DAT + second spray with zinc sulphate @ 3 % on 30 DAT

T₉ - ZSS + First spray with zinc sulphate @ 3 % on 15 DAT + second spray with potassium silicate @ 3 % on 30 DAT

T₁₀ - Carbendazim (0.1 %) – Comparison

T₁₁ - Un treated control.

Phenolic changes - Method of sampling

Samples of plant materials from each treatment were taken at 0, 7, 14 and 21 days after inoculation both in healthy and inoculated plants for estimating the changes in the biochemical constituents viz., reducing sugars, non-reducing sugars, total sugars, starch, ortho dihydroxy phenols, total phenols, amino nitrogen, protein and enzymes like peroxidase, polyphenol oxidase, phenylalanine ammonia lyase and ascorbic acid oxidase.

Preparation of ethanol extracts [6]

Plant materials of both healthy and infected were collected and 4 g samples were taken. They were chopped and then extracted in 16 ml of boiling 80 per cent ethanol for 5 min. and cooled in running tap water. The material was homogenized by grinding in a porcelain pestle and mortar and squeezed through two layers of cheese cloth. The residue was transferred back to 5 ml of boiling 80 per cent ethanol and reextracted for 5 min. cooled and filtered. Both the extracts were pooled and filtered through Whatman No.41 filter paper. A jet of ethyl alcohol was used to wash the filter paper and the final volume was adjusted to 20 ml with 80 per cent ethyl alcohol, so as to get 5 ml of the extract representing every g of plant tissue. The ethanol extract was used for the estimation of sugars, phenols, amino nitrogen and protein. The biochemical constituents were assessed based on standard procedures.

Biochemical constituents	References
Amino nitrogen	Moore and Stein, 1958 [7]
Starch	Summer and Somers, 1949 [8]
Total Protein	Lowry <i>et al.</i> 1951 [9]

3. RESULTS AND DISCUSSION

Post infectional biochemical changes

Changes in Amino nitrogen

Application of ZS, SA and PS altered the amino nitrogen content in rice variety ADT 36. Application of SA at 15 DAT or its combination with ZS, and PS decreased the amino nitrogen content. The maximum content of 7.77 mg was recorded in control at the end of the sampling period. The minimum content of 4.19 mg was observed in T₆ (ZSS + SA₁ + PS₂). Amino nitrogen content was gradually increased up to 14th day of sampling and then decreased to the initial level in T₆ treatment on 21st day of sampling. Application of SA at 15 DAT reduced the amino nitrogen content than application of SA at 30 DAT (Table 1).

Table 1. Changes in Amino nitrogen of rice var. ADT 36 as influenced by application of ZS, SA, PS and *B. oryzae* inoculation

T.No.	Treatments	Amino nitrogen (mg/g)			
		0 (days)	7 (days)	14 (days)	21 (days)
1	ZSS + ZSF ₁ + ZSF ₂	5.01	5.92	9.07	4.92
2	ZSS + SA ₁ + SA ₂	4.52	5.19	8.43	4.44
3	ZSS + PS ₁ + PS ₂	4.93	5.73	8.91	4.76
4	ZSS + ZSF ₁ + SA ₂	4.46	5.14	8.31	4.40
5	ZSS + SA ₁ + ZSF ₂	4.40	5.06	8.23	4.34
6	ZSS + SA ₁ + PS ₂	4.34	4.98	8.12	4.19
7	ZSS + PS ₁ + SA ₂	4.57	5.32	8.56	4.60
8	ZSS + PS ₁ + ZSF ₂	4.80	5.67	8.86	4.69
9	ZSS + ZSF ₁ + PS ₂	4.69	5.42	8.67	4.57
10	Carbendazim	5.82	6.82	9.89	5.48
11	Control	7.92	8.55	11.89	7.77

Changes in Protein

Protein content was significantly altered by ZS, SA and PS application (Table 2). Application of SA either on 15 DAT or on 30 DAT or in combination with PS increased the protein content in ADT 36. The maximum level of protein was observed in T₆ with 55.82 mg/g at the initial sampling, whereas at the final sampling it was increased to the maximum (66.48 mg/g). Protein content increased up to the 14th day of sampling and then slightly reduced in all the treatments. The minimum protein content was recorded in control at final sampling (40.62 mg/g).

Table 2. Changes in Protein content of rice var. ADT 36 as influenced by application of ZS, SA, PS and *B. oryzae* inoculation

T. No.	Treatments	Protein content (mg/g)			
		0 (days)	7 (days)	14 (days)	21 (days)
1	ZSS + ZSF ₁ + ZSF ₂	52.04	56.92	58.61	53.19
2	ZSS + SA ₁ + SA ₂	53.51	57.86	59.66	54.83
3	ZSS + PS ₁ + PS ₂	52.35	57.05	58.66	53.27
4	ZSS + ZSF ₁ + SA ₂	52.70	57.23	58.80	53.64
5	ZSS + SA ₁ + ZSF ₂	53.63	58.12	59.92	54.09
6	ZSS + SA ₁ + PS ₂	55.82	61.93	70.56	66.48
7	ZSS + PS ₁ + SA ₂	54.14	59.06	62.16	57.81

8	ZSS + PS ₁ + ZSF ₂	53.37	57.61	59.45	55.06
9	ZSS + ZSF ₁ + PS ₂	53.12	57.49	59.23	55.00
10	Carbendazim	43.90	47.28	48.33	44.86
11	Control	36.29	40.08	43.88	40.62

Starch

ZS, SA and PS application significantly influenced the starch content in rice variety ADT 36 when compared to control. Maximum content of starch was observed in T₆ (ZSS + SA₁ + PS₂) which recorded 53.15 mg/g at the 14th day of the sampling. In control, the starch content was the least with 33.35 mg at the end of the sampling period. Starch content gradually increased up to 14th day of sampling and then slightly decreased in all the treatments. The treatment combinations with SA recorded significant increase in starch content (Table 3).

Table 3. Changes in Starch content of rice var. ADT 36 as influenced by application of ZS, SA, PS and *B. oryzae* inoculation

T. No.	Treatments	Starch content (mg/g)			
		0 (days)	7 (days)	14 (days)	21 (days)
1	ZSS + ZSF ₁ + ZSF ₂	34.94	42.49	47.13	44.91
2	ZSS + SA ₁ + SA ₂	36.27	45.43	49.82	48.12
3	ZSS + PS ₁ + PS ₂	35.05	42.82	47.66	45.63
4	ZSS + ZSF ₁ + SA ₂	36.69	45.68	52.74	50.86
5	ZSS + SA ₁ + ZSF ₂	37.05	45.93	52.87	51.04
6	ZSS + SA ₁ + PS ₂	37.48	46.13	53.15	51.80
7	ZSS + PS ₁ + SA ₂	37.25	46.04	52.98	51.36
8	ZSS + PS ₁ + ZSF ₂	35.17	43.31	48.26	46.87
9	ZSS + ZSF ₁ + PS ₂	35.89	43.68	48.99	47.00
10	Carbendazim	33.60	37.23	42.58	39.61
11	Control	31.64	32.42	35.99	33.35

Treatment combinations with ZnSO₄, salicylic acid and potassium silicate decreased the amino nitrogen content and increased the protein content in rice variety ADT 36 (Table 1 and 2). Nitrogenous compounds are important for the growth and multiplication of invading pathogens. Miswa and Miyasakai [10] observed that bacterial leaf blight susceptible rice variety contained greater amount of amino nitrogen than the resistant variety. Close correlation between blast susceptibility and soluble nitrogen was reported by Chen [11]. Karpagavalli and Ramabadran [12] reported reduction in amino nitrogen content due to lignite fly ash application. Ragavan [13] and Jaiganesh [14] stated reduction in amino

nitrogen content due to chemical inducers like salicylic acid and nicotinic acid application. Involvement of protein components in plant disease resistance has been documented in several plant patho-systems [15], [16], [17]. Increased protein content due to salicylic acid application in rust infected groundnut leaves [18] and blast affected rice [14] has also been reported. Nutrient management is an integral component of efficient production agriculture. Nutrient sufficiency may provide a general form of disease resistance by maintaining a high level of inhibitory compounds in tissue or quick response to invasion by a pathogen [19]. The inorganic salts like silicates, bicarbonates, phosphates, chlorides and phosphites have properties that make them desirable for inclusion in Integrated Disease Management (IDM) programmes [20]. Their proposed advantages include low cost, favourable safety profile for humans and the environment and low mammalian toxicity [21], [22]. Silicon can reduce the impact of nutrient deficiency of phosphorus in crop plants [23] and can reduce the toxicity effects caused by aluminium and iron [24] and induce defense responses [25]. Thus, silicon fertilizer could be an environmental friendly alternative to manage crop diseases [26]. Some economically important diseases in rice such as blast, brown spot, stem rot and grain discoloration have been reduced by silicon nutrient application with increased production [27], [28], [29], [30]. These reports lend support to the present findings.

4. CONCLUSION

Among the various integrated disease management approaches, Soil Application of Zinc sulphate @ 25 kg/ha and foliar application of Salicylic acid @ 50 ppm on 15 Days after transplanting and Foliar application of Potassium silicate @ 3 % recorded the minimum disease incidence and its was proved that the positive effect of Biochemical changes.

ACKNOWLEDGEMENT

The authors are greatly thankful for Annamalai University authorities for given the wonderful research facilities

CONFLICT OF INTEREST

Authors have no any conflict of interest.

REFERENCES

1. Van Nguyen N, Ferrero A and Meeting the challenges of global rice production. Paddy water Environ., 2006; 4: 1-9.
2. Anonymous A and “Top Rice Producing Countries”; International Trade Commodities (www.Internationaltradecommodities.com) 2008.
3. Tamil Nadu Statistics Report and Department of Economics and statistics, Government of India, Season, Area and Crop production report 2006-2007. 2007 (Ref.: www.tn.gov.in)
4. Ou S H and Rice Diseases, 2nd Edition, Common Wealth Mycological Institute, U.K. 1985; 380p.

5. Sridhar R, Nayak N and Physiology of disease resistance in rice. In : Basic Research for Crop Disease Management (Edn. By P. Vidhyasekaran), Daya Publishing House, New Delhi, India. 1990;110-120.
6. Sridhar R, Mahadevan A and Triggering mechanism in rice blast disease. Acta Phytopath. Acad. Sci. Hung, 1968; 3: 415-421.
7. Moore S, Stein W H and The modified reagent for the photometrical determination of amino acids and related compounds J.Biol. Chem. 1958; 211: 907-913.
8. Summer J B, Somers G F and Laboratory Experiments in Biological Chemistry, 2nd ed. Academic Press, Inc., New York, 1949; 173 p.
9. Lowry O H, Rosebrough, N J, Farr, A L, Randall R J and Protein measurement with Folin-phenol reagent. J.Biol. Chem., 1951; 153: 263-265.
10. Miswa T, Miyasakai E and Studies on the leaf blight of rice plant. 1. Alteration of content of carbohydrates, nitrogenous and phosphorus compounds in the diseased leaves. Annu. Phytopath. Soc. Japan, 1972; 38: 375-380.
11. Chen Z Q and Investigations on the mechanism of resistance to rice blast disease. J. South China Agricultural University. 1989;10: 82-91.
12. Karpagavalli S, Ramabadran R and Influence of lignite fly-ash on the biochemical changes in rice towards disease resistance. Proceedings of National Seminar on Fly ash Utilization, held at Neyveli Lignite Corporation, Neyveli, India on 22 & 23 Aug. 1996, 146-156 pp.
13. Ragavan R and Studies on the management of blast disease of paddy incited by *Pyricularia oryzae* Cavara. Ph.D. Thesis , Annamalai University, Tamil nadu , 2003.
14. Jaiganesh V and Studies on the management of rice blast caused by *Pyricularia oryzae* Cavara. M.Sc.(Ag.), Thesis, Annamalai University, India, 2005.
15. Carvalho M F, Turgeon R, Lazarowitz S G and The geminivirus nuclear shuttle protein NSP inhibits the activity of AtNSI, a vascular-expressed *Arabidopsis* acetyltransferase regulated with the sink-to-source transition. Plant Physiol., 2006; 140: 1317–1330
16. Aboshosha S S, Atta Alla, S I, EL- Korany A E, EL- Argawy E and Protein analysis and peroxidase isozymes as molecular markers for resistance and susceptibility of sunflower to *Macrophomina phaseolina*. International Journal of Agriculture and Biology, 2008; 10 (1): 28-34.
17. Mishra B K, Mishra RK, Mishra R C, Tiwari A K, Yadav R S, Dikshit A and Biocontrol efficacy of *Trichoderma viride* isolates against fungal pathogens causing disease in *Vigna radiate* L. Archives of applied science research, 2011; 3(2): 361-369.
18. Siddaramaiah A L, Desai S A, Hegde R K and Studies on estimation of loss due to rust and late leaf spot of groundnut. Mysore Journal of Agricultural Sciences, 1983;17: 365-367.
19. Huber D M, Haneklaus S and Managing nutrition to control plant diseases. A review.

20. Deliopoulos T, Kettlewell P S, Hare M C and Fungal disease suppression by inorganic salts. *Crop Protection*, 2010; 29: 1059-1075.
21. Reuveni M, Reuveni R and Efficacy of foliar sprays of Phosphates in controlling Powdery Mildews in field-grown nectarine, mango trees and grapevines. *Crop Protection*, 1995; 14:311-314.
22. Olivier C, MacNeil C R, Loria R and Application of organic and inorganic salts to field-grown potato tubers can suppress silver scurf during potato storage. *Plant Dis.*, 1999; 83: 814-818.
23. Epstein E and The anomaly of silicon in plant biology. *Proceedings of the National Academy of Sciences of the United States of America*, 1994; 91: 11-17.
24. Marschner H and *Mineral Nutrition of Higher Plants*, 2nd edn. London, UK, Academic Press, 1995; p. 889.
25. Fawe A, Abou-Zaid M, Menzies J G, Belanger R R and Silicon-mediated accumulation of flavonoid phytoalexins in cucumber. *Phytopathol.*, 1998; 88: 396–401.
26. Cai K, Gao D, Luo S, Zeng R, Yang J, Zhu X and Physiological and cytological mechanisms of silicon induced resistance in rice against blast disease. *Physiologia Planta.*, 2008; 134: 324-333.
27. Seebold K W, Datnoff L E, Correa-Victoria F J, Kucharek T A, Snyder G H and Effect of silicon rate and host resistance on blast, scald and yield of upland rice. *Plant Dis.*, 2000; 84: 871–876.
28. Aziz T, Gill M A, Rahmatullah R and Silicon nutrition and crop production: A review. *Pak. J. Agric. Sci.*, 2002; 39 (3):181-187.
29. Zhang Z Y, Jin Y, Huang C J, Duan Y Q, Yang Y H, Yang H Q and Effect of phosphorus and Zinc nutrition on tobacco mosaic virus. *J. Hu. Agri. Univ.*, 2008; DOI: CNKI:SUN:HNND.0.2008-03-014
30. Santos G R D, Neto M D de, Ramos L N, Sarmento R A, Korndorfer G H, Ignacio M and Effect of silicon source on rice diseases and yield in the state of Tocantins, Brazil. *Acta Scientiarum Agronomy*, 2011; 33: 451-456.